


# Innovations in Incidence Geometry


Algebraic, Topological and Combinatorial



**More configurations on elliptic curves**

Marco Bramato, Lorenz Halbeisen and Norbert Hungerbühler





## More configurations on elliptic curves

Marco Bramato, Lorenz Halbeisen and Norbert Hungerbühler

*Dedicated to the memory of Branko Grünbaum*

We construct elliptic  $(3r_s, sr_3)$  configurations for all integers  $r \geq s \geq 1$ . This solves an open problem of Branko Grünbaum. The configurations which we build have mirror symmetry and even  $D_3$  symmetry if  $r$  is a multiple of 3. The configurations are dynamic in the sense that the points can be moved along the elliptic curve in such a way that all line incidences are preserved.

### 1. Introduction

**1.1. Elliptic configurations.** The study of configurations has a long and rich history. We refer to Grünbaum [3] as a main reference, and the bibliography in [7] for an overview of newer developments. To fix the notation, let  $p, l, \pi, \lambda \in \mathbb{N}$ . Then a  $(p_\lambda, l_\pi)$  configuration is a set of  $p$  points and  $l$  lines in the projective plane such that each point is incident to  $\lambda$  lines and each line is incident to  $\pi$  points. If  $p = l$  and consequently  $\lambda = \pi$ , we just write  $(p_\lambda)$  instead of  $(p_\lambda, l_\pi)$ .

Of particular interest are *elliptic configurations*, i.e., configurations whose points lie on an elliptic curve or, more specifically, in the torsion group of an elliptic curve. A long standing open problem is the question of Grünbaum [3, Section 4.8, Open problem 4]: for which integers  $r$  are there elliptic  $(3r_4, 4r_3)$  configurations? Notice that for  $r = 3$ , the Hesse configuration  $(9_4, 12_3)$  can be realized in the complex projective plane as the set of inflection points of an elliptic curve, but it has no realization with straight lines in the Euclidean or projective plane because of the Sylvester–Gallai theorem. On the other hand, examples of elliptic  $(12_4, 16_3)$  configurations were found by Grünbaum [3, p. 249], Coxeter [1, p. 440], and Feld [2] (where one can find also an example of an elliptic  $(36_7, 84_3)$  configuration). For an elliptic  $(24_6, 48_3)$  configuration, see [6]. In [11], Metelka identified eight elliptic

---

MSC2020: 51A05, 51A20.

Keywords: configurations, elliptic curves.

(12<sub>4</sub>, 16<sub>3</sub>) configurations. Elliptic curves carrying Pappos, Desargues or Fano configurations have been discussed in [10].

Recently, some progress was reported in [7], where elliptic  $((p-1)_3)$  configurations are constructed for every prime  $p > 7$ . There are  $(3r_4, 4r_3)$  configurations whenever  $3r = p - 1$  for some prime  $p > 7$ , and for every  $k \geq 2$  there is an elliptic  $(9k_4, 12k_3)$  configuration with  $D_3$  symmetry (the symmetry group of an equilateral triangle).

In the present article we generalize the ideas in [7] and show that elliptic  $(3r_4, 4r_3)$  configurations exist for all  $r \geq 4$  and that one can even construct elliptic  $(3r_s, sr_3)$  configurations for any  $s \in \mathbb{N}$  and  $r \geq s$ . We offer constructions with  $D_1$  symmetry (mirror symmetry), and, if  $r$  is a multiple of 3,  $D_3$  symmetry. A particularly pleasing property is that the configurations are dynamic in the sense that one can move the points of the configurations along the elliptic curve in such a way that all line incidences and rotational symmetries are preserved (see [9] for recent results in this context). Thus, Grünbaum's question is completely answered.

**1.2. Elliptic curves.** Recall that an elliptic curve in Weierstrass normal form

$$\Gamma_{g_2, g_3} : y^2 = 4x^3 - g_2x - g_3$$

in  $\mathbb{C}^2$  can be parametrized by the Weierstrass function  $\wp(z) := \wp(z, g_2, g_3)$  via

$$\gamma : \mathbb{C} \rightarrow \bar{\mathbb{C}}^2, \quad z \mapsto (\wp(z), \wp'(z)).$$

For real  $g_2, g_3$ , the Weierstrass function has a real period  $\omega_1$  and an imaginary period  $\omega_2$ . Then, depending on the parameters  $g_2, g_3$ , the curve  $\Gamma_{g_2, g_3}$  in  $\mathbb{R}^2$  consists of one or two connected components. We will call the unbounded component *the odd branch*, and the bounded component *the even branch*. The odd branch is parametrized by

$$\gamma_{\text{odd}} : (0, \omega_1) \rightarrow \mathbb{R}^2, \quad t \mapsto (\wp(t), \wp'(t)),$$

and the even branch, if it exists, by

$$\gamma_{\text{even}} : [0, \omega_1) \rightarrow \mathbb{R}^2, \quad t \mapsto \left( \wp\left(t + \frac{1}{2}\omega_2\right), \wp'\left(t + \frac{1}{2}\omega_2\right) \right)$$

(see [1, p. 441]). The projective version of the curve  $\Gamma_{g_2, g_3}$  carries the group operation  $\oplus$  of an elliptic curve with the neutral element  $\mathcal{O} = (0, 1, 0)$  which corresponds to  $\gamma(0)$ . The group operation is compatible with the parametrization, i.e.,

$$\gamma(z_1) \oplus \gamma(z_2) = \gamma(z_1 + z_2) \tag{1}$$

(see [12, Chapter VI, §3] for the general theory of elliptic curves, and [8, §3] for the parametrization by the Weierstrass function).

Every regular cubic curve can be brought, by a projective transformation, into the  $D_3$  symmetric normal form

$$\mathbb{C}_{D_3} : x^3 - 3xy^2 - 3(b-3)(x^2 + y^2) + 4b^2(b-9) = 0$$

with  $b \in \mathbb{R} \setminus \{1\}$  (see [7]). Depending on the parameter  $b$ , the curve  $\mathbb{C}_{D_3}$  consists only of the odd branch or of both the odd and the even branch (see Section 2.1). One of the three symmetry axes is the  $x$ -axis, and the rotational  $C_3$  symmetry is with respect to the origin.

The configurations which we want to build will be constructed with the help of the arithmetic structure of  $\Gamma_{g_2, g_3}$  and  $\Gamma_{D_3}$ , respectively. Let  $Q := \gamma((q\omega_1)/n)$  for some integer parameter  $q$  with  $(q, n) = 1$ . By (1),  $Q$  has order  $n$  and generates a group  $G_n$  which is isomorphic to  $\mathbb{Z}/n\mathbb{Z}$ . To fix the notation, let

$$P_u := u * Q = \underbrace{Q \oplus \cdots \oplus Q}_{u \text{ times}} = \gamma\left(\frac{uq\omega_1}{n}\right),$$

and we can identify the point  $P_u$  on the curve with  $u \in \mathbb{Z}/n\mathbb{Z}$ . If we take, in addition, the points  $P'_u := \gamma((uq\omega_1)/n + \omega_2/2)$ , we obtain a group isomorphic to  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ : the points  $P_u$  correspond to  $\{0\} \times \mathbb{Z}/n\mathbb{Z}$ , and the points  $P'_u$  to  $\{1\} \times \mathbb{Z}/n\mathbb{Z}$ . In particular, we have

$$\begin{aligned} P'_{u_1} \oplus P'_{u_2} &= \gamma\left(\frac{qu_1\omega_1}{n} + \frac{\omega_2}{2}\right) \oplus \gamma\left(\frac{qu_2\omega_1}{n} + \frac{\omega_2}{2}\right) \\ &= \gamma\left(\frac{q(u_1 + u_2)\omega_1}{n} + \omega_2\right) = \gamma\left(\frac{q(u_1 + u_2)\omega_1}{n}\right) = P_{u_1 + u_2}. \end{aligned}$$

**1.3. Notation.** We identify points  $P_u$  on the odd branch of the curve directly with the corresponding value  $(0, u)$  in  $\{0\} \times \mathbb{Z}/n\mathbb{Z}$ , and similarly, we identify  $(1, u)$  with points  $P'_u$  on the even branch. We will also call the former *odd points* and the latter *even points*. When speaking of any point  $u$  (without specifying the branch), we mean a point on any component of the curve or the corresponding element in  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ . Finally, instead of writing  $\oplus$  we just write  $+$  from now on.

If the points  $u, v, w$  are collinear, then the corresponding line is denoted by  $[u, v, w]$ . Three distinct points  $u, v, w$  on the elliptic curve are collinear if and only if their sum (interpreted as sum on the elliptic curve or in  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ ) is zero, and we just write  $u + v + w = 0$ .

Our configurations will only consist of lines that contain 3 points, and hence are of the form  $[(0, u), (0, v), (0, w)]$  or  $[(0, u), (1, v), (1, w)]$  (where the order of points could be changed). We call the former *0-lines* and the latter *1-lines*.

**1.4. Operations on lines.** We introduce three operations on lines: a conjugation, a rotation, and a translation.

If  $[u, v, w]$  is a line, then  $-[u, v, w] := [-u, -v, -w]$  is the *conjugate line* of  $[u, v, w]$ . In the context of a configuration on a  $\Gamma_{D_3}$  curve, the line and the conjugate line are mirror symmetric with respect to the  $x$ -axis. Hence, an elliptic configuration on a  $\Gamma_{D_3}$  curve is mirror symmetric with respect to the  $x$ -axis if each of its conjugate lines is an element of the configuration.

Let  $k \in \mathbb{Z}$ . For any point  $u$  in  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$  we define

$$\rho_k(u) := u + (0, k).$$

If  $[u, v, w]$  is a line with points in  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3k\mathbb{Z}$  for some positive integer  $k$ , we can define the *rotated line*

$$\rho_k([u, v, w]) := [\rho_k(u), \rho_k(v), \rho_k(w)].$$

Note that this indeed yields a line, since  $\rho_k(u) + \rho_k(v) + \rho_k(w) = u + v + w + (0, 3k)$ . In the context of a configuration on a  $\Gamma_{D_3}$  curve, this rotation corresponds to a rotation with angle  $2\pi/3$  about the origin. So, an elliptic configuration on a  $\Gamma_{D_3}$  curve has rotational  $C_3$  symmetry with respect to the origin if each of its rotated lines is an element of the configuration.

If  $S_t = \gamma_{\text{odd}}(t)$  is a point on the odd branch of an elliptic curve and  $\varepsilon$  is an arbitrary real number, then the *translated point*  $S_t^\varepsilon$  is defined by  $S_t^\varepsilon := \gamma_{\text{odd}}(t - 2\varepsilon)$ . Similarly, if  $S'_t = \gamma_{\text{even}}(t)$  is a point on the even branch, the translated point is defined by  $S'_t{}^\varepsilon := \gamma_{\text{even}}(t + \varepsilon)$ . If the three points  $S_{t_1}, S'_{t_2}, S'_{t_3}$  form a 1-line, then the translated 1-line is given by the points  $S_{t_1}^\varepsilon, S'_{t_2}{}^\varepsilon, S'_{t_3}{}^\varepsilon$ . Indeed these three points are on a line, since

$$\begin{aligned} S_{t_1}^\varepsilon \oplus S'_{t_2}{}^\varepsilon \oplus S'_{t_3}{}^\varepsilon &= \gamma(t_1 - 2\varepsilon) \oplus \gamma\left(t_2 + \varepsilon + \frac{\omega_2}{2}\right) \oplus \gamma\left(t_3 + \varepsilon + \frac{\omega_2}{2}\right) \\ &= \gamma(t_1 + t_2 + t_3 + \omega_2) = S_{t_1} \oplus S'_{t_2} \oplus S'_{t_3} = 0. \end{aligned}$$

## 2. Simple elliptic $(3r_s, sr_3)$ configurations

**Definition 1.** Let  $r$  and  $s$  be positive integers. A *simple elliptic*  $(3r_s, sr_3)$  configuration has  $r$  points on the odd branch and  $2r$  points on the even branch.

The reason for the name “simple” is that a simple configuration only consists of 1-lines: Indeed, if there are  $l_0$  0-lines and  $l_1$  1-lines, then the number of odd points is  $3l_0 + l_1$  while the number of even points is  $2l_1$ . So, for a simple configuration we have

$$2l_1 = 2(3l_0 + l_1) \Rightarrow l_0 = 0.$$

We will also see that simple configurations are given by quite simple constructions.

**Definition 2.** An elliptic configuration is called *dynamic* if one can move the points along the curve in such a way that all line incidences are preserved.

In particular, an elliptic configuration is dynamic if every point of the curve is a point of a configuration of the same type.

**Lemma 3.** *An elliptic configuration which only consists of 1-lines is dynamic.*

*Proof.* We can apply for an arbitrary real  $\varepsilon$  the corresponding translation as defined in Section 1.4 simultaneously to all points of the configuration. Then, every 1-line is translated to the 1-line through the translated points.  $\square$

Below, we will prove the following result.

**Theorem 4.** *For all  $r \geq s \geq 1$  there exists a simple  $(3r_s, sr_3)$  configuration with  $D_3$  symmetry if  $r \equiv 0 \pmod{3}$  and  $D_1$  symmetry otherwise. These configurations are dynamic. The translated versions of  $D_3$  symmetric configurations have rotational  $C_3$  symmetry.*

We first consider the case when  $s = 4$ .

**2.1. Construction of simple  $(3r_4, 4r_3)$  configurations.** We start by constructing elliptic  $(3r_4, 4r_3)$  configurations for  $r \geq 4$ . If  $3 \mid r$ , then these configurations will have  $D_3$  symmetry.

Let  $r \geq 4$ . The following points are understood to be in  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2r\mathbb{Z}$ . The set of configuration points is  $\mathcal{P} := \mathcal{P}_0 \cup \mathcal{P}_1$ , where

$$\begin{aligned}\mathcal{P}_0 &:= \{(0, 1), (0, 3), (0, 5), \dots, (0, 2r - 1)\}, \\ \mathcal{P}_1 &:= \{(1, 0), (1, 1), (1, 2), \dots, (1, 2r - 1)\}.\end{aligned}$$

$\mathcal{P}$  contains  $r + 2r = 3r$  points as needed. As mentioned previously, we only use 1-lines. Define  $\mathcal{L}_1$  to be the set of the lines

$$\begin{aligned}[a_0, b_0, c_0] &:= [(0, -1), (1, 0), (1, 1)], \\ [a_{i+1}, b_{i+1}, c_{i+1}] &:= [a_i - (0, 2), b_i + (0, 1), c_i + (0, 1)] \quad \text{for } i \in \{0, \dots, 2r - 2\}\end{aligned}$$

and let  $\mathcal{L}_2$  be the set of the lines


$$\begin{aligned}[d_0, e_0, f_0] &:= [(0, -3), (1, 0), (1, 3)], \\ [d_{i+1}, e_{i+1}, f_{i+1}] &:= [d_i - (0, 2), e_i + (0, 1), f_i + (0, 1)] \quad \text{for } i \in \{0, \dots, 2r - 2\}.\end{aligned}$$

Observe that

$$\begin{aligned}[a_i, b_i, c_i] &= [(0, -1 - 2i), (1, i), (1, 1 + i)], \\ [d_i, e_i, f_i] &= [(0, -3 - 2i), (1, i), (1, 3 + i)],\end{aligned}$$

where all numbers are to be read modulo  $2r$ . We claim that  $\mathcal{L}_1 \cup \mathcal{L}_2$  yields appropriate lines for a  $(3r_4, 4r_3)$  configuration. This is seen by considering the following points:

- The elements of  $\mathcal{L}_1 \cup \mathcal{L}_2$  define 1-lines in  $\mathcal{P}$ : All  $a_i$  and  $d_i$  are odd points and lie in  $\mathcal{P}$ . All  $b_i, c_i, e_i, f_i$  are even points and lie in  $\mathcal{P}$ . Furthermore,  $a_i + b_i + c_i = 0 = d_i + e_i + f_i$  and no two points of a line are equal.



**Figure 1.** Simple  $D_3$  symmetric  $(18_4, 24_3)$  configuration, with parameter  $q = 1$ . Adding the twelve lines  $(0, \pm 1 + 4k)$ ,  $(1, \pm 2 + 4k)$ ,  $(1, \pm 9 + 4k)$  and  $(0, \pm 1 + 4k)$ ,  $(1, \pm 3 + 4k)$ ,  $(1, \pm 8 + 4k)$  for  $k = 1, 2, 3$ , we can extend this to a  $(18_6, 36_3)$  configuration. This is a special case of the configurations treated in Section 2.2.

- All lines are different: If  $a_i = a_j$  for  $i \neq j$ , then the sets  $\{b_i, c_i\}$  and  $\{b_j, c_j\}$  are different. So, all lines in  $\mathcal{L}_1$  are different. Similarly, all lines in  $\mathcal{L}_2$  are different. Moreover, the lines in  $\mathcal{L}_1$  are different from those in  $\mathcal{L}_2$  since the differences of the even points on a line are  $(0, 1)$  and  $(0, 3)$ , respectively. Note that for  $r = 3$  we would obtain two equal lines since then  $[d_0, e_0, f_0] = [(0, -3), (1, 0), (1, 3)] = [d_3, f_3, e_3]$ .
- Each point in  $\mathcal{P}_0$  occurs twice in the set of the  $a_i$  and twice in the set of the  $d_i$ . And the lines of both  $\mathcal{L}_1$  and  $\mathcal{L}_2$  contain each point in  $\mathcal{P}_1$  twice.


Concerning the symmetry of the configuration, observe the following: For  $i \in \mathbb{Z}$ ,

$$-[a_i, b_i, c_i] = [(0, 2i + 1), (1, -i), (1, -i - 1)] = [a_{-i-1}, c_{-i-1}, b_{-i-1}]$$

taking all indices modulo  $2r$ . Therefore the lines in  $\mathcal{L}_1$  form a  $D_1$  symmetry. Similarly, this can be seen to be true for the lines in  $\mathcal{L}_2$ , where  $-[d_i, e_i, f_i] \equiv [d_{-i-3}, f_{-i-3}, e_{-i-3}]$  for all  $i \in \mathbb{Z}$ .

This completes the proof of Theorem 4 for  $s = 4$ . Figure 1 shows the smallest simple  $D_3$  symmetric configuration according to this construction, while Figure 2





**Figure 2.** A translated version of the simple  $D_3$  symmetric  $(184, 24_3)$  configuration in Figure 1.


is a translated version of it. Figure 3 shows a simple  $D_1$  symmetric  $(33_4, 44_3)$  configuration.

We even obtain a  $D_3$  symmetry for these configurations if  $r$  is a multiple of 3. To see this, let  $n := \frac{2r}{3}$  and notice that since  $n \equiv -2n \pmod{2r}$  we have

$$\begin{aligned} \rho_n[a_i, b_i, c_i] &= [(0, n - 2i - 1), (1, i + n), (1, i + n + 1)] = [a_{i+n}, b_{i+n}, c_{i+n}], \\ \rho_n[d_i, e_i, f_i] &= [(0, n - 2i - 3), (1, i + n), (1, i + n + 3)] = [d_{i+n}, e_{i+n}, f_{i+n}]. \end{aligned}$$

Since the configuration is simple, it follows from Lemma 3 that it is dynamic. Also observe that if three points on  $\Gamma_{D_3}$  are vertices of an equilateral triangle centered at the origin, then the translated points have also this property. So, when we translate a  $D_3$  symmetric elliptic configuration on a  $\Gamma_{D_3}$  curve, there results an elliptic configuration with rotational  $C_3$  symmetry.

**2.2. Simple  $(3r_s, sr_3)$  configurations for arbitrary  $r \geq s \geq 1$ .** Now, we generalize the construction of the previous section to  $(3r_s, sr_3)$  configurations where  $r \geq s \geq 1$ .



**Figure 3.** Simple  $D_1$  symmetric  $(33_4, 44_3)$  configuration, with parameter  $q = 5$ .

For the proof we have to distinguish two cases which we consider separately in the following subsections. The first subsection covers the case  $r \geq s \geq 1$  with  $r$  odd or  $r = s$  even, the second subsections treats the case  $r > s \geq 1$  with  $r$  even.

**2.2.1.  $r$  odd or  $r = s$  even.** Let  $r \geq s \geq 1$  with  $r$  odd or  $r = s$  even. We use the same set of points  $\mathcal{P} := \mathcal{P}_0 \cup \mathcal{P}_1$  as in Section 2.1. For  $1 \leq j \leq \lceil \frac{s-1}{2} \rceil =: \bar{s}$  let  $\mathcal{L}^j$  be the set consisting of the lines

$$[a_0^j, b_0^j, c_0^j] := [(0, 1-2j), (1, 0), (1, 2j-1)],$$

$$[a_{i+1}^j, b_{i+1}^j, c_{i+1}^j] := [a_i^j - (0, 2), b_i^j + (0, 1), c_i^j + (0, 1)] \quad \text{for } i \in \{0, \dots, 2r-2\},$$

and  $\mathcal{L}^*$  the set consisting of the lines

$$[d_0, e_0, f_0] := [(0, -r), (1, 0), (1, r)],$$

$$[d_{i+1}, e_{i+1}, f_{i+1}] := [d_i - (0, 2), e_i + (0, 1), f_i + (0, 1)] \quad \text{for } i \in \{0, \dots, r-2\}.$$

Observe that

$$[a_i^j, b_i^j, c_i^j] = [(0, 1-2j-2i), (1, i), (1, 2j+i-1)],$$

$$[d_i, e_i, f_i] = [(0, -r-2i), (1, i), (1, r+i)].$$

When  $s$  is even, we claim that  $\mathcal{L} := \bigcup_{j=1}^{\bar{s}} \mathcal{L}^j$  yields appropriate lines for a  $(3r_s, sr_3)$  configuration, and if  $s$  is odd, then  $\mathcal{L} := \mathcal{L}^* \cup \bigcup_{j=1}^{\bar{s}} \mathcal{L}^j$  does. To see this, we check the following:

- (1) The elements of  $\mathcal{L}$  define 1-lines in  $\mathcal{P}$ : All  $a_i^j$  and  $d_i$  are odd and lie in  $\mathcal{P}$  and so do the even points  $b_i^j, c_i^j, e_i^j$  and  $f_i^j$ . Furthermore,  $a_i^j + b_i^j + c_i^j = 0 = d_i + e_i + f_i$  for all  $i, j$ , and no two points of a line are equal.
- (2) All lines are different: For different  $j$ , the difference between  $b_i^j$  and  $c_i^j$  is different, and different from the difference between  $e_i$  and  $f_i$ . For fixed  $j$  all sets  $\{b_i^j, c_i^j\}$  are different. The same can be checked for the sets  $\{e_i, f_i\}$ .
- (3) If  $s$  is odd, each point in  $\mathcal{P}_0$  occurs  $s - 1$  times in the set of the  $a_i^j$  and once in the set of the  $d_i$ . The lines of all  $\mathcal{L}^j$  and  $\mathcal{L}^*$  together contain each point in  $\mathcal{P}_1$  exactly  $s$  times. Similarly, when  $s$  is even, each point in  $\mathcal{P}$  is contained in  $s$  lines.

To see the  $D_1$  symmetry, observe that

$$\begin{aligned} -[a_i^j, b_i^j, c_i^j] &= [(0, 2i + 2j - 1), (1, -i), (1, 1 - i - 2j)] \\ &= [a_{1-2j-i}^j, c_{1-2j-i}^j, b_{1-2j-i}^j], \end{aligned}$$

where indices are taken modulo  $2r$ . Hence the lines in  $\mathcal{L}^j$  are  $D_1$  symmetric. Similarly, we have  $-[d_i, e_i, f_i] = [d_{r-i}, f_{r-i}, e_{r-i}]$ .

To see that  $D_3$  symmetry occurs when  $r$  is a multiple of 3, let  $n := \frac{2r}{3}$  and note that  $\rho_n[a_i^j, b_i^j, c_i^j] = [(0, 1 - 2j - 2i + n), (1, i + n), (1, i + n + 2j - 1)] = [a_{i+n}^j, b_{i+n}^j, c_{i+n}^j]$  and

$$\rho_n[d_i, e_i, f_i] = [(0, -r - 2i - 2n), (1, i + n), (1, i + n + r)] = [d_{i+n}, e_{i+n}, f_{i+n}].$$

Finally, since also in the present construction only 1-lines are used, the  $(3r_s, sr_3)$  configurations are dynamic.

**2.2.2.** *r even and  $r > s \geq 1$ .* Let  $r > s \geq 1$  with  $r$  even and define

$$\mathcal{P}_0 := \{(0, 0), (0, 2), \dots, (0, 2r - 2)\} \quad \text{and} \quad \mathcal{P}_1 := \{(1, 0), (1, 1), \dots, (1, 2r - 1)\}$$

and  $\mathcal{P} = \mathcal{P}_0 \cup \mathcal{P}_1$ . For  $1 \leq j \leq \lceil \frac{s-1}{2} \rceil =: \bar{s}$  let  $\mathcal{L}^j$  be the set consisting of the lines


$$[a_0^j, b_0^j, c_0^j] := [(0, -2j), (1, 0), (1, 2j)],$$

$$[a_{i+1}^j, b_{i+1}^j, c_{i+1}^j] := [a_i^j - (0, 2), b_i^j + (0, 1), c_i^j + (0, 1)] \quad \text{for } i \in \{0, \dots, 2r - 2\},$$

and  $\mathcal{L}^*$  be the set consisting of the lines

$$[d_0, e_0, f_0] := [(0, -r), (1, 0), (1, r)],$$

$$[d_{i+1}, e_{i+1}, f_{i+1}] := [d_i - (0, 2), e_i + (0, 1), f_i + (0, 1)] \quad \text{for } i \in \{0, \dots, r - 2\}.$$



**Figure 4.** A  $D_3$  symmetric  $(365, 60_3)$  configuration, with parameter  $q = 5$ .

Observe that

$$\begin{aligned} [a_i^j, b_i^j, c_i^j] &= [(0, -2j - 2i), (1, i), (1, 2j + i)], \\ [d_i, e_i, f_i] &= [(0, -r - 2i), (1, i), (1, r + i)]. \end{aligned}$$

It can be checked very similarly as in the previous section that  $\mathcal{L} := \bigcup_{j=1}^s \mathcal{L}^j$  if  $s$  is even, and  $\mathcal{L} := \mathcal{L}^* \cup \bigcup_{j=1}^s \mathcal{L}^j$  if  $s$  is odd, yield appropriate lines for a  $(3r_s, rs_3)$  configuration. Again we obtain  $D_1$  symmetry for all such configurations and  $D_3$  symmetry for  $r \equiv 0 \pmod{3}$ .

Note that here we use the point  $(0, 0)$  at infinity in the  $D_1$  symmetric case, and the points  $(0, 0)$ ,  $(0, \frac{2r}{3})$  and  $(0, \frac{4r}{3})$  at infinity in the  $D_3$  symmetric case. Since the configuration is dynamic, we can still obtain a  $D_3$  symmetric configuration with all points finite if we choose  $\varepsilon = (3\omega_1)/(4r)$  for the translation. Figure 4 shows such a situation for  $r = 12$ ,  $s = 5$  and parameter  $q = 5$ .

### 3. Closing remarks and open problems

For a fixed curve  $\Gamma_{D_3}$  and fixed numbers  $r \geq s \geq 1$  there is still some freedom in our construction of a  $(3r_s, sr_3)$  configuration: we can choose the parameter  $\varepsilon \in \mathbb{R}$  (see Section 1.4) and the parameter  $q \in \mathbb{N}$  (see Section 1.2). However the resulting configurations are combinatorially isomorphic: there is a bijection of the points which preserves the line incidences. But choosing a different  $\varepsilon$  leads in general to configurations which are geometrically nonisomorphic: there is no projective map which maps one configuration to the other. Indeed, the configuration in Figure 4 with  $\varepsilon = (3\omega_1)/(4r)$  cannot be geometrically isomorphic to the corresponding configuration with  $\varepsilon = 0$ , since the latter occupies the three inflection points of the curve (at infinity), while the former does not. Also the choice of different values for the parameter  $q$  leads in general to geometrically nonisomorphic configurations, since a projective map preserves the order of the points on both branches of the curve, while different values of  $q$  lead in general to a different order of the points (compare Figures 1 and 3).

We conclude with two open problems.

- Does every  $(3r_s, sr_3)$  configuration of the combinatorial type constructed in Section 2 have its points necessarily on a cubic curve?
- Which curves other than cubic curves carry dynamic configurations?

Observe that the miraculous chains of Poncelet polygons introduced in [4] and [5] are examples of dynamic configurations carried by conics. So far, only a  $(9_3)$  configuration consisting of three Poncelet triangles and a  $(24_3)$  configuration consisting of six Poncelet quadrilaterals are known.

### References

- [1] H. S. M. Coxeter, “Self-dual configurations and regular graphs”, *Bull. Amer. Math. Soc.* **56** (1950), 413–455. MR Zbl
- [2] J. M. Feld, “Configurations inscriptible in a plane cubic curve”, *Amer. Math. Monthly* **43:9** (1936), 549–555. MR Zbl
- [3] B. Grünbaum, *Configurations of points and lines*, Graduate Studies in Mathematics **103**, American Mathematical Society, Providence, RI, 2009. MR Zbl
- [4] L. Halbeisen and N. Hungerbühler, “Conjugate conics and closed chains of Poncelet polygons”, *Mitt. Math. Ges. Hamburg* **36** (2016), 5–28. MR Zbl
- [5] L. Halbeisen and N. Hungerbühler, “Closed chains of conics carrying Poncelet triangles”, *Beitr. Algebra Geom.* **58:2** (2017), 277–302. MR Zbl
- [6] L. Halbeisen and N. Hungerbühler, “Constructing cubic curves with involutions”, *Beitr. Algebra Geom.* **63:4** (2022), 921–940. MR Zbl
- [7] A. Halbeisen, L. Halbeisen, and N. Hungerbühler, “Configurations on elliptic curves”, *Innov. Incidence Geom.* **19:3** (2022), 111–135. MR Zbl

- [8] G. A. Jones and D. Singerman, *Complex functions: an algebraic and geometric viewpoint*, Cambridge University Press, 1987. MR Zbl
- [9] S. Lundqvist, K. Stokes, and L.-D. Öhman, “Exploring the rigidity of planar configurations of points and rods”, *Discrete Appl. Math.* **336** (2023), 68–82. MR
- [10] N. S. Mendelsohn, R. Padmanabhan, and B. Wolk, “Placement of the Desargues configuration on a cubic curve”, *Geom. Dedicata* **40**:2 (1991), 165–170. MR Zbl
- [11] V. Metelka, “Über ebene Konfigurationen  $(12_4, 16_3)$ , die mit einer irreduziblen Kurve dritter Ordnung inzidieren”, *Časopis Pěst. Mat.* **91** (1966), 261–307. MR Zbl
- [12] J. H. Silverman, *The arithmetic of elliptic curves*, 2nd ed., Graduate Texts in Mathematics **106**, Springer, 2009. MR Zbl

Received 26 Jan 2023.

MARCO BRAMATO:

bramatom@student.ethz.ch

Department of Mathematics, ETH Zürich, Zürich, Switzerland

LORENZ HALBEISEN:

lorenz.halbeisen@math.ethz.ch

Department of Mathematics, ETH Zürich, Zürich, Switzerland

NORBERT HUNGERBÜHLER:

norbert.hungerbuehler@math.ethz.ch

Department of Mathematics, ETH Zürich, Zürich, Switzerland

# Innovations in Incidence Geometry

msp.org/iig

## MANAGING EDITORS

Anton Betten	Colorado State University anton.betten@colostate.edu
Michael Cuntz	Leibniz Universität Hannover cuntz@maths.uni-hannover.de
Dimitri Leemans	Université Libre de Bruxelles leemans.dimitri@ulb.be
Oliver Lorscheid	University of Groningen o.lorscheid@rug.nl
James Parkinson	University of Sydney jamesp@maths.usyd.edu.au
Koen Thas	Ghent University koen.thas@gmail.com

## EDITORS

Takuro Abe	Kyushu University
Philippe Cara	Vrije Universiteit Brussel
Antonio Cossidente	Università della Basilicata
Hans Cuypers	Eindhoven University of Technology
Bart De Bruyn	Ghent University
Tom De Medts	Ghent University
Alice Devillers	University of Western Australia
Maria Elisa Fernandes	University of Aveiro
Massimo Giulietti	Università degli Studi di Perugia
James Hirschfeld	University of Sussex
Alessandro Montinaro	Università di Salento
Gabi Nebe	RWTH Aachen University
Piotr Pokora	Pedagogical University of Cracow
Anna Pun	University of Virginia
Bertrand Rémy	École Polytechnique
Felipe Rincón	Queen Mary University of London
Binod Sahoo	NISER, Bhubaneswar
Jeroen Schillewaert	Auckland University
Petra Schwer	University of Magdeburg
Kris Shaw	University of Oslo
Klara Stokes	Umea Universitet
Joseph A. Thas	Ghent University
Hendrik Van Maldeghem	Ghent University

## PRODUCTION

Silvio Levy	(Scientific Editor) production@msp.org
-------------	---

---

See inside back cover or [msp.org/iig](http://msp.org/iig) for submission instructions.

The subscription price for 2024 is US \$265/year for the electronic version, and \$315/year (+\$10, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Innovations in Incidence Geometry: Algebraic, Topological and Combinatorial (ISSN 2640-7345 electronic, 2640-7337 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

---

IIG peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY  
 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>  
© 2024 Mathematical Sciences Publishers

# Innovations in Incidence Geometry

Vol. 21 No. 1

2024

Frobenius actions on Del Pezzo surfaces of degree 2	1
OLOF BERGVALL	
A note on the rank 5 polytopes of $M_{24}$	11
VERONICA KELSEY, ROBERT NICOLAIDES and PETER ROWLEY	
Victoris–Rips complexes of platonic solids	17
NADA SALEH, THOMAS TITZ MITE and STEFAN WITZEL	
An incomplete real tree with complete segments	35
RAPHAEL APPENZELLER, LUCA DE ROSA, XENIA FLAMM and VICTOR JAECK	
More configurations on elliptic curves	45
MARCO BRAMATO, LORENZ HALBEISEN and NORBERT HUNGERBÜHLER	
Flag-accurate arrangements	57
PAUL MÜCKSCH, GERHARD RÖHRLE and TAN NHAT TRAN	
The generalized Sylvester’s and orchard problems via discriminantal arrangement	117
PRAGNYA DAS, ELISA PALEZZATO and SIMONA SETTEPANELLA	
Locally resolvable BIBDs and generalized quadrangles with ovoids	131
RYAN MCCULLOCH	

